

NARAYANA'S SENSATIONAL SUCCESS ACROSS INDIA

7 Students Secured **100 Percentile**
in All India JEE Main-2020

BELOW 10
21
RANKS
All Cat

BELOW 100
113
RANKS
All Cat



ADMISSIONS OPEN (2020-21)

OUR REGULAR CLASSROOM PROGRAMME

One Year Classroom Program
JEE/NEET-2021
(for students moving from XI to XII)

Two Year Classroom Program
JEE/NEET-2022
(for students moving from X to XI)

Three Year Integrated Classroom Program
JEE/NEET-2023
(for students moving from IX to X)

Four Year Integrated Classroom Program
JEE/NEET-2024
(for students moving from VIII to IX)

FOUNDATION PROGRAMMES
For NTSE, NSEJS, JSTSE,
Olympiads & School/Board Exams
(for students moving to
Class VI, VII, VIII, IX & X)

APEX BATCH
Two years school Integrated
Classroom Program - 2022
For JEE Main & Advance / NEET (for XI Studying Students)
Course Feature - Complete Coverage of CBSE - Regular Classes - Weekly Test & Regular Analysis - Lab Facility
- Motivation & Counseling - Competitive Exam Prep - Ample time for self study

Online Classes for IIT/NEET/Foundation/Olympiads

- Access Recording of Past Classes on n-Learn App
- Online Parent Teacher Meeting
- Personalized Extra Classes & Live Doubt Solving
- Hybrid/Customized Classroom model
- Video Solution of Weekly/Fortnightly Test
- Printed Study Material will be sent by us
- n-Learn App
- Counselling Motivational sessions
- Affordable Fee
- Doubt Classes / Practice Classes
- Provision to Convert from online to regular classroom programme
- Once Classes resume by just paying nominal fee

Online Test

- Micro & Macro Analysis
- Relative performance (All India Ranking)
- Question wise Analysis
- Unlimited Practice Test
- Grand Test

NARAYANA

Digital
Classes
STUDY ONLINE FROM HOME

For Class
7th to 12th +



JEE-MAIN-2021

MARCH ATTEMPT

17.03.21_SHIFT - II

CHEMISTRY

CHEMISTRY

1. Match the followings-

- | | |
|------------------------------------|-----------------------|
| (A) Artificial sweetner | (i) Sodium benzoate |
| (B) Antiseptic | (ii) Bithional |
| (C) Preservative | (iii) Sodium stearate |
| (D) Glyceryl ester of stearic acid | (iv) Sucralose |

(1) (A) → (iv), (B) → (ii), (C) → (i), (D) → (iii)

(2) (A) → (iii), (B) → (i), (C) → (ii), (D) → (iv)

(3) (A) → (i), (B) → (iii), (C) → (i), (D) → (iii)


(4) (A) → (i), (B) → (iii), (C) → (iii), (D) → (i)

Ans. (1)

2. Kjeldahl method is applicable for

- (1) PhN_2^{\oplus} (2) Ph-NO_2 (3) $\text{Ph-CH}_2\text{-NH}_2$ (4) 

Ans. (3)



Which of the following enzymes are used in above reactions respectively?

- | | |
|--------------------------|---------------------------|
| (1) Amylase and Zymase | (2) Invertase and Zymase |
| (3) Zymase and Invertase | (4) Amylase and Invertase |

Ans. (2)

4. Fructose is an example of

- (1) Pyranose (2) Aldohexose (3) Ketohexose (4) Pentose

Ans. (3)

5. **Statement-1** : 2-Methylbutane is oxidised by KMnO_4 to give 2-Methyl butan-2-ol.

Statement-2 : An alkane is easily oxidised by KMnO_4 .

- (1) Both Statement-1 and Statement-2 are correct
- (2) Both Statement-1 and Statement-2 are false
- (3) Statement-1 is correct and Statement-2 is false
- (4) Statement-1 is false and Statement-2 is correct

Ans. (3)



6. 1° , 2° and 3° amines can be distinguish by-

- (1) Chloroform and KOH
- (2) CS_2 with HgCl_2
- (3) Tosyl chloride
- (4) $\text{HCl} + \text{ZnCl}_2$

Ans. (3)

7. How many carbon-carbon σ bonds are present in mesityl oxide ?

Ans. (5)




Correct statement about A & B is -

- (1) A is more stable and formed with faster rate.
- (2) B is more stable and formed with faster rate.
- (3) A is less stable and formed with slow rate.
- (4) B is less stable and formed with faster rate.


Ans. (1)

9. FeCl_3 is reacted with oxalic acid in presence of KOH. Find secondary valency of iron in product

Ans. (6)





Secondary valency = 6



Identify A, B and C

	A	B	C
(1)		CH_3CHO	HCl
(2)		CH_3CHO	HCl
(3)			HCl
(4)			HCl

Ans. (2)



Ans. (77)

Sol. Moles of $\text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} = \frac{140.5}{140.5} = 1 \text{ mol.}$

Moles of $\text{Ph}-\overset{\text{O}}{\parallel}{\text{C}}-\text{N}(\text{Ph})_2$ that should be obtained by mol-mol analysis = 1 mol

Theoretical mass of product = $1 \times 273\text{g}$

Observed mass of product = 210 g

Percentage yield of product = $\frac{W_{\text{experimental}}}{W_{\text{theoretical}}} \times 100 = \frac{210}{273} \times 100 = 76.9\%$

Ans. 77

12. Element with atomic number 24 is expected to show following common oxidation states -

- (1) +1 to +6 (2) +1 & +3 to +6
 (3) +3 to +6 (4) +2 to +6

Ans. (4)

13. Match the column-

- (A) $[\text{Cu}(\text{NH}_3)_4][\text{CuCl}_4]$ (P) Solvate isomerism
 (B) $[\text{Co}(\text{H}_2\text{O})_6]\text{Cl}_3$ (Q) Coordination isomerism
 (C) $[\text{Co}(\text{NH}_3)_3\text{Cl}_3]$ (R) Optical isomerism
 (D) $\text{Cis}-[\text{Co}(\text{en})_2\text{Cl}_2]^+$ (S) Geometrical isomerism

(1) $\text{A} \rightarrow \text{Q}, \text{B} \rightarrow \text{P}, \text{C} \rightarrow \text{S}, \text{D} \rightarrow \text{R}$

(2) $\text{A} \rightarrow \text{P}, \text{B} \rightarrow \text{Q}, \text{C} \rightarrow \text{S}, \text{D} \rightarrow \text{R}$

(3) $\text{A} \rightarrow \text{P}, \text{B} \rightarrow \text{Q}, \text{C} \rightarrow \text{R}, \text{D} \rightarrow \text{S}$

(4) $\text{A} \rightarrow \text{S}, \text{B} \rightarrow \text{R}, \text{C} \rightarrow \text{P}, \text{D} \rightarrow \text{Q}$

Ans. (1)

14. Match the following ores with their chemical formula :

- (A) Bauxite (P) $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}$
 (B) Haematite (Q) Fe_2O_3
 (C) Magnetite (R) Fe_3O_4
 (D) Malachite (S) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$

(1) $\text{A} \rightarrow \text{P} ; \text{B} \rightarrow \text{Q} ; \text{C} \rightarrow \text{R} ; \text{D} \rightarrow \text{S}$

(2) $\text{A} \rightarrow \text{S} ; \text{B} \rightarrow \text{R} ; \text{C} \rightarrow \text{Q} ; \text{D} \rightarrow \text{P}$

(3) $\text{A} \rightarrow \text{R} ; \text{B} \rightarrow \text{P} ; \text{C} \rightarrow \text{S} ; \text{D} \rightarrow \text{Q}$

(4) $\text{A} \rightarrow \text{P} ; \text{B} \rightarrow \text{Q} ; \text{C} \rightarrow \text{S} ; \text{D} \rightarrow \text{R}$

Ans. (1)

Ans. (354)

Sol. $K_P = K_C (RT)^1$
 $600.1 = 20.4 (0.083T)$
 $T \approx 354 \text{ K}$

- 16.** 1 molal aqueous $\text{K}_4[\text{Fe}(\text{CN})_6]$ having $\alpha = 0.4$ has same boiling point as 18.1% by weight solution of non electrolyte A. Find molar mass of A.

Ans. (85)


Sol. Since B.P. is same \Rightarrow elevation in B.P. is also same for both solution

$$\begin{aligned}(\Delta T_B)_{K_4[Fe(CN)_6]} &= (\Delta T_B)_A \\ \Rightarrow (ik_b m)_{K_4[Fe(CN)_6]} &= (ik_b m)_A \\ &= (1 + 4\alpha) \times 1 = 1 \times \frac{(18.1) / M \times 1000}{(100 - 18.1)} \\ \Rightarrow 2.6 &= \frac{(18.1)}{M} \times \frac{1000}{(81.9)} \Rightarrow M = 85\end{aligned}$$

- 17.** Linear species is:


- (1) N_3^- (2) NO_2 (3) Cl_2O (4) O_3

Ans. (1)




- 18.** In which of the following process entropy of system is decreasing?

- (A) Freezing of water at 0°C (B) Freezing of water at -10°C
(C) Adsorption of H_2 on Pb (D) Dissolution of NaCl in H_2O



- (1) A, B, C, E (2) A, B, C, D (3) A, B, C, D, E (4) A, B

Ans. (1)



Remaining (A), (B), (C) and (E) have negative entropy

19. $2A + B_2 \rightarrow 2AB$ is an elementary reaction. If volume of container is reduced to $\frac{1}{3}$ rd. Determine ratio of rate final to initial.

Ans. (27)

Sol. For elementary reaction,

$$\text{Rate of reaction} = K [A]^2 [B_2]$$

$$\text{Initial rate} = K \left(\frac{n_A}{v_0} \right)^2 \left(\frac{n_B}{v_0} \right)$$

$$\text{Final rate} = K \left(\frac{n_A}{\frac{v_0}{3}} \right)^2 \left(\frac{n_B}{\frac{v_0}{3}} \right) = 27 K \left(\frac{n_A}{v_0} \right)^2 \left(\frac{n_B}{v_0} \right) \Rightarrow \frac{\text{Final Rate}}{\text{Initial Rate}} = \frac{27}{1}$$

20. Spin only magnetic moment in ground state of iron is $x \times 10^{-1}$.

$$(\sqrt{2} = 1.41, \sqrt{3} = 1.73)$$

Ans. (49)

Sol. $\text{Fe} - 1s^2 2s^2 2p^6 3s^2 3p^6 3d^6 4s^2$

Number of unpaired electron = 4

$$N_{\text{spin}} = \sqrt{n(n+2)}$$

$$= \sqrt{4(4+2)}$$

$$= \sqrt{24}$$

$$= 4.9$$

$$= 49 \times 10^{-1}$$

21. A conductivity cell when filled with NaCl solution is found to have conductivity $0.14 \Omega^{-1} \text{m}^{-1}$ and $R = 4.09 \Omega$. When HCl solution is filled in same conductivity cell, R is found to be 1.03Ω . If conductivity of HCl solution is $x \times 10^{-2}$ (in $\Omega^{-1} \text{m}^{-1}$). Determine 'x'.

Ans. (56)

Sol. for NaCl solution

$$R = \left(\frac{1}{K} \right) \left(\frac{\ell}{A} \right) \Rightarrow \frac{\ell}{A} = (R)(K) = (4.09)(0.14) \text{ m}^{-1}$$

for HCl solution

$$R = \left(\frac{1}{K} \right) \left(\frac{\ell}{A} \right) \Rightarrow K = \frac{\left(\frac{\ell}{A} \right)}{R} = \frac{(4.09)(0.14)}{1.03} = 56 \times 10^{-2}$$

$$x = 56$$

22. Number of atoms in 20 ml of Cl_2 at STP are $x \times 10^{21}$. Find x

$$R = 0.083$$

$$N_A = 6.023 \times 10^{23}$$

Ans. (1)

Sol. $n = \frac{PV}{RT}$

$$= \frac{1 \times 20 \times 10^{-3}}{0.083 \times 273}$$

$$\text{Number of atoms} = \frac{1 \times 20 \times 10^{-3}}{0.083 \times 273} \times 2 \times 6.023 \times 10^{23}$$

$$= 1.06 \times 10^{21}$$

Ans.1

23. If NaCl is doped with 10^{-3} mole percentage of SrCl_2 , cationic vacancies per mole of NaCl. ($N_A = 6.023 \times 10^{23}$) are 6.022×10^x . Determine x.

Ans. (18)

Sol. 100 mole NaCl \longrightarrow 10^{-3} mole $\text{SrCl}_2 \longrightarrow 10^{-3} N_A$ Cationic vacancies

\therefore 1 mole NaCl $\longrightarrow 10^{-5} N_A$ Cationic vacancies

$$= 10^{-5} \times 6.023 \times 10^{23}$$

$$= 6.022 \times 10^{18} \text{ Cationic vacancies}$$

24. During the recovery of NH_3 in solvey process byproduct formed is :

(1) CaCl_2

(2) Na_2CO_3

(3) NaCl

(4) Ca(OH)_2

Ans. (1)

25. Highest flocculating power for the coagulation of negatively charged sol is –

(1) Na^+

(2) Be^{2+}

(3) PO_4^{3-}

(4) SO_4^{2-}

Ans. (2)