

# NARAYANA'S SENSATIONAL SUCCESS ACROSS INDIA

## 7 Students Secured 100 Percentile in All India JEE Main-2020

BELOW 10

**21**  
RANKS

All Cat

BELOW 100

**113**  
RANKS

All Cat

M LIKHITH REDDY HT No. TL0305724	SHIVA KRISHNA HT No. TL01330751	TANUJA HT No. TL01327005	SWAYAM CHUBE HT No. MR16303069	HARSHVARDHAN AGARWAL HT No. HR09300789	YESHASH CHANDRA HT No. TL02305081	PARTH DWIVEDI AppL No. 200310166791
-------------------------------------	------------------------------------	-----------------------------	-----------------------------------	---	--------------------------------------	--

## ADMISSIONS OPEN (2020-21)

### OUR REGULAR CLASSROOM PROGRAMME

**One Year Classroom Program**  
**JEE/NEET-2021**  
(for students moving from XI to XII)

**Two Year Classroom Program**  
**JEE/NEET-2022**  
(for students moving from X to XI)

**Three Year Integrated Classroom Program**  
**JEE/NEET-2023**  
(for students moving from IX to X)

**Four Year Integrated Classroom Program**  
**JEE/NEET-2024**  
(for students moving from VIII to IX)

**FOUNDATION PROGRAMMES**  
**For NTSE, NSEJS, JSTSE,**  
**Olympiads & School/Board Exams**  
(for students moving to  
Class VI, VII, VIII, IX & X)

**APEX BATCH**  
**Two years school Integrated**  
**Classroom Program - 2022**  
For JEE Main & Advance / NEET (for XI Studying Students)  
**Course Feature** - Complete Coverage of CBSE - Regular Classes - Weekly Test & Regular Analysis - Lab Facility  
- Motivation & Counseling - Competitive Exam Prep - Ample time for self study

**Online Classes for IIT/NEET/Foundation/Olympiads**

- Access Recording of Past Classes on n-Learn App
- Online Parent Teacher Meeting
- Personalized Extra Classes & Live Doubt Solving
- Hybrid/Customized Classroom model
- Video Solution of Weekly/Fortnightly Test
- Printed Study Material will be sent by us
- n-Learn App
- Counselling Motivational sessions
- Affordable Fee
- Doubt Classes / Practice Classes
- Provision to Convert from online to regular classroom programme
- Once Classes resume by just paying nominal fee

**Online Test**

- Micro & Macro Analysis
- Relative performance (All India Ranking)
- Question wise Analysis
- Unlimited Practice Test
- Grand Test

**NARAYANA**

**Digital**  
**Classes**  
STUDY ONLINE FROM HOME

**For Class**  
**7<sup>th</sup> to 12<sup>th</sup> +**





**JEE-MAIN-2021**

**18.03.21\_SHIFT - I**

**MARCH ATTEMPT**

**THE NARAYANA GROUP**

**PHYSICS**

## JEE(MAIN) 2021 (18 MARCH ATTEMPT) SHIFT-1

### PHYSICS

1. If a simple pendulum completes 200 oscillation in 100 sec. Least count of watch is 1 sec., length of simple pendulum is 100 cm and it's least count is 1 mm then find max. possible percentage error in measuring acceleration due to gravity.

- (1) 3.2                      (2) 5.2                      (3) 2.1                      (4) 4.1

Ans. (3)

Sol.  $T = 2\pi\sqrt{\frac{\ell}{g}}$

$$T^2 = 4\pi^2 \left( \frac{\ell}{g} \right)$$

$$g = 4\pi^2 \left( \frac{\ell}{T^2} \right)$$

$$\frac{\Delta g}{g} = \frac{\Delta \ell}{\ell} + 2 \frac{\Delta T}{T}$$

$$\frac{\Delta g}{g} \times 100 = \frac{0.1\text{cm}}{100\text{cm}} \times 100\% + 2 \left( \frac{1\text{sec}}{100\text{sec}} \right) \times 100\%$$


$$\frac{\Delta g}{g} \times 100 = 2.1\%$$

2. A girl is looking at the distant rectangular window, she finds window to be blurred & non-uniformly curved. What eye defect she may have?

- (1) Myopia & Astigmatism                      (2) Myopia & Hypermetropia  
 (3) Astigmatism                      (4) Hypermetropia & Astigmatism

Ans. (1)

3. In the circuit shown evaluate potential difference across  $10\Omega$  in volts?



Ans. 70.00

**Sol.**  $R_{eq} = 10 + \frac{10 \times 20}{50 \times 20}$

$$= \frac{170}{7} \Omega$$

$$I = \frac{V}{R_{eq}} = \frac{170}{170} \times 7 = 7 \text{ amp}$$

$$V_{10\Omega} = IR$$

$$= 7 \times 10 = 70 \text{ v}$$

4. A satellite revolves in a circular orbit of radius R around earth with time period T. Find its time period if it starts revolving in radius 9R?

- (1) 3T                      (2) 6T                      (3) 9T                      (4) 27T

**Ans.** (4)


**Sol.**  $T^2 \propto R^3$

$$\therefore \left( \frac{T_2}{T_1} \right)^2 = \left( \frac{R_2}{R_1} \right)^3$$

$$\therefore \left( \frac{T_2}{T_1} \right)^2 = 9^3$$

$$\therefore \frac{T_2}{T_1} = 27$$

5. AC circuit diagram is shown. Find time taken to reach its current from  $i_{rms}$  to  $i_{max}$ .



- (1) 10 milli sec.                      (2) 1 milli sec.                      (3) 2.5 milli sec.                      (4) 5 milli sec.

**Ans.** (3)

**Sol.**  $i = i_{max} \sin(\omega t + \theta)$

at  $t = 0$ ,  $i = i_{rms}$

$$i_{rms} = \sqrt{2} (i_{rms}) \sin \theta$$

$$\theta = \frac{\pi}{4}$$



$$i = i_{\max} \sin \left( \omega t + \frac{\pi}{4} \right)$$

$$\text{at } t = t_1, i = i_{\max}$$

$$i_{\max} = i_{\max} \sin \left( \omega t_1 + \frac{\pi}{4} \right)$$

$$\omega t_1 + \frac{\pi}{4} = \frac{\pi}{2}$$

$$\omega t_1 + \frac{\pi}{4}$$

$$\frac{2\pi}{T} t_1 = \frac{\pi}{4}$$

$$t_1 = \frac{T}{8}$$

$$t_1 = \frac{1}{8} \left( \frac{1}{f} \right) = \frac{1}{8} \left( \frac{1}{50} \right)$$

$$t_1 = \frac{1000}{400} \text{ m sec} = 2.5 \text{ m sec}$$

6. In LCR circuit L and C are constant and R is increased then:
- (1) Quality factor and resonant frequency both are unchanged.
  - (2) Quality factor is increased.
  - (3) Band width is increased.
  - (4) Quality factor remains unchanged

**Ans.** (3)

**Sol.**  $\omega = \frac{1}{\sqrt{LC}}, Q = \frac{1}{R} \sqrt{\frac{L}{C}}, \text{ Band width} = \frac{R}{L}$

7. In YDSE setup, distance between slits is 0.5 mm & separation between slits plane & screen is 0.5 m. Find the distance between 1<sup>st</sup> maxima & 3<sup>rd</sup> maxima if light used has wave length 5890 Å.
- (1)  $1178 \times 10^{-6} \text{ m}$       (2)  $1178 \times 10^{-7} \text{ m}$       (3)  $1178 \times 10^{-8} \text{ m}$       (4)  $5890 \times 10^{-7} \text{ m}$

**Ans.** (1)

**Sol.** Distance between 1<sup>st</sup> & 3<sup>rd</sup> maxima will be  $3\beta$ .

$$\therefore 2 \times \frac{\lambda D}{d} = 2 \times 5890 \times 10^{-10} \times \frac{0.5}{0.5 \times 10^{-3}}$$

$$= 11780 \times 10^{-7} \text{ m}$$

8. A closed current carrying loop is placed in uniform magnetic field. Then in equilibrium shape of wire will be :
- (1) straight
  - (2) unchanged
  - (3) circular and plane perpendicular to magnetic field
  - (4) Circular and plane parallel to magnetic field

Ans. (3)

9. A muon particle (mass =  $207 m_e$ ) revolves around hydrogen nucleus. Find its ionisation energy.?  
[ $m_e$  = mass of electron]

- (1) 13.6 eV                      (2) 27.2 eV                      (3)  $13.6 \times 207$  eV                      (4) 331.8 eV

Ans. (Bonus)


Sol.  $E_n = -13.6 \times \frac{\mu}{m_e} \text{ eV}$

$$\mu = \frac{(1836m_e)(207m_e)}{(1836 + 207)m_e}$$

$$= \frac{1836 \times 207}{2043} = 186 m_e.$$

$$\therefore \text{Ionisation energy} = 13.6 \times 186 \text{ eV}$$

10. An object is moving with constant acceleration. Choose the correct option.



Ans. (4)



**Sol.**  $a = \text{constant}$

$$v \propto t$$

$$x \propto t^2$$

- 11.** A ring of mass  $M$  is rotating with constant angular velocity  $\omega$  about axis of rotation passing through centre and perpendicular to the plane of ring. Two particles each of mass  $m$  are placed gently diametrically at opposite position. Find new angular velocity.

(1)  $\left(\frac{M+2m}{M}\right)\omega$       (2)  $\left(\frac{M\omega}{M+2m}\right)$       (3)  $\left(\frac{M-2m}{M}\right)\omega$       (4)  $\left(\frac{m\omega}{M+2m}\right)$

**Ans.** (3)

**Sol.** Using angular momentum conservation

$$L_i = MR^2\omega$$

$$L_f = (MR^2 + 2mR^2)\omega'$$

$$\omega' = \left(\frac{M\omega}{M+2m}\right)$$

- 12.** Electromagnetic wave is propagating in  $x$  direction. Magnetic field in space is given by  $\vec{B} = 2 \times 10^{-8}(\text{T})\hat{k}$ . What will be the value and direction of electric field.

(1)  $0.6 \hat{j}$       (2)  $6 \hat{j}$       (3)  $0.6 \hat{k}$       (4)  $6 \hat{k}$

**Ans.** (2)

**Sol.**  $E = CB$

$$E = 3 \times 10^8 \times 2 \times 10^{-8}$$

$$E = 6$$

direction of  $\vec{v}$  is  $\vec{E} \times \vec{B}$

$$\hat{i} = \hat{j} \times \hat{k}$$

$$\text{so } \vec{E} = 6\hat{j}$$

- 13.** A machine starting from Rest delivers constant Power 'P'. Then distance travelled by it in time 't' is proportional to:-

(1)  $t^{-3/2}$       (2)  $t^{1/2}$       (3)  $t^{3/2}$       (4)  $t^{-1/2}$

**Ans.** (3)

**Sol.**  $P = Fv$

$$P = mav$$

$$P \int dt = m \int v dv$$

$$m \frac{v^2}{2} = Pt$$

$$v = \left( \frac{2Pt}{m} \right)^{1/2}$$

$$\frac{dx}{dt} = \left( \frac{2Pt}{m} \right)^{1/2}$$

$$x = \left( \frac{2P}{m} \right)^{1/2} \frac{t^{3/2}}{\frac{3}{2}}$$

$$x \propto t^{3/2}$$

14. An object is performing SHM with time period 2 sec. If time taken by it to move from mean position to half of amplitude is  $\frac{1}{K}$  sec. Then value of K is.

- (1) 3                      (2) 6                      (3) 4                      (4) 2


**Ans.** (2)

**Sol.** from 0 to  $\frac{A}{2}$

$$\text{time} = \frac{T}{12} \text{ sec}$$

$$\frac{2}{12} = \frac{1}{6} \text{ sec}$$

15. In given P-V graph process CA is adiabatic. Find work done in process CA if gas is diatomic ( $\gamma = 1.4$ ):




- (1) -400 J                      (2) -500 J                      (3) 200                      (4) 400

**Ans.** (2)

**Sol.**  $W = \frac{nR\Delta T}{1-\gamma} = \frac{P_2 V_2 - P_1 V_1}{1-\gamma} = \frac{200 \times 3 - 100 \times 4}{1-1.4} = -500 \text{ J}$



16. Four identical solenoids are connected as shown in figure



If magnetic field in A is 3T, evaluate magnetic field in C

- (1) 1T (2) 9T (3) 12T (4) 6T

Ans. (1)

Sol.  $B_A = \mu_0 n I = 3T$

$$B_C = \mu_0 n \frac{I}{3}$$

$$B_C = 1T$$

17. In a wire  $V = 5.0V$ ,  $I = 2.00A$ ,  $L = 10.0$  cm and diameter  $d = 5.00$  mm. Evaluate  $\frac{\Delta\rho}{\rho} \times 100$  ?

- (1) 3.9% (2) 1.9% (3) 2.9% (4) 3%

Ans. (1)

Sol.  $\frac{\Delta\rho}{\rho} = \frac{\Delta R}{R} + \frac{\Delta\ell}{\ell} + \frac{2\Delta d}{d}$

$$\frac{\Delta\rho}{\rho} = \frac{\Delta V}{V} + \frac{\Delta I}{I} + \frac{\Delta\ell}{\ell} + \frac{2\Delta d}{d}$$

$$\frac{\Delta\rho}{\rho} \% = \left( \frac{0.1}{5} + \frac{0.01}{2} + \frac{0.1}{10} + 2 \times \frac{0.01}{5} \right) \times 100$$

$$= 2 + 0.5 + 1 + 0.4 = 3.9\%$$

18. A is forming B and C independently if  $A \rightarrow B$  with half life =  $T_{1/2}(B)$  and if  $A \rightarrow C$  with half life  $T_{1/2}(C)$  then what will be overall half life:

(1)  $\frac{T_{1/2}(B) \times T_{1/2}(C)}{T_{1/2}(B) + T_{1/2}(C)}$

(2)  $\frac{T_{1/2}(B) + T_{1/2}(C)}{T_{1/2}(B) \times T_{1/2}(C)}$

(3)  $\frac{T_{1/2}(B) \times T_{1/2}(C)}{T_{1/2}(B) - T_{1/2}(C)}$

(4)  $\frac{T_{1/2}(B) + T_{1/2}(C)}{T_{1/2}(B) - T_{1/2}(C)}$

Ans. (1)

**Sol.**  $-\frac{dN_A}{dt} = \lambda_B N_A + \lambda_C N_A$

$$= (\lambda_B + \lambda_C) N_A = \lambda_{eq} N_A$$

$$\lambda_{eq} = \lambda_B + \lambda_C$$

$$\frac{\ln 2}{T_{eq}} = \frac{\ln 2}{T_{1/2B}} + \frac{\ln 2}{T_{1/2C}} \Rightarrow \frac{1}{T_{eq}} = \frac{1}{T_{1/2B}} + \frac{1}{T_{1/2C}}$$

$$T_{eq} = \frac{T_{1/2B} \times T_{1/2C}}{T_{1/2B} + T_{1/2C}}$$

- 19.** Two wires A and B of same material having elongation 2 mm and 4 mm respectively on applying 2N take. If radius of B is four times the radius of A and ratio of length of A is to B in the form of  $\frac{1}{x}$  then the value of x is

**Ans.** 32.00

**Sol.**  $\frac{F}{A} = Y \frac{\Delta L}{L}$

$$\frac{F}{\pi r_A^2} = Y \frac{\Delta L_A}{L_A} \quad \text{----- (i)}$$

$$\frac{F}{\pi r_B^2} = Y \frac{\Delta L_B}{L_B} \quad \text{----- (ii)}$$

$$\left(\frac{r_B}{r_A}\right)^2 = \frac{\Delta L_A}{\Delta L_B} \times \frac{L_A}{L_B} \quad r_B = 4r_A$$

$$16 = \frac{2}{4} \times \frac{L_B}{L_A} \quad \frac{r_B}{r_A} = 4$$

$$\frac{L_B}{L_A} = 32$$

$$\frac{a}{b} = \frac{1}{32}$$


$$x = 32$$

- 20.** A man is swimming in a river at an angle  $120^\circ$  with river flow. Speed of man in still water is 10m/s. If he reaches the other bank exactly opposite to origin point, find speed of flow of river (in m/s)

**Ans.** 5.00



**Sol.**



Net speed perpendicular to line.

AB must be zero.

$$\therefore v_m \sin 30 = V_R$$

$$\therefore v_r = 5 \text{ m/s}$$

- 21.** If ratio of de-Broglie wavelength of particle and electron is 2 : 1 and ratio of their velocity is 4 : 1. Then

- (1) mass of particle is 8 times that of electron
- (2) mass of electron is 8 times that of particle
- (3) mass of electron is 16 times that of particle
- (4) mass of particle is 16 times that of electron

**Ans.** (2)

**Sol.**

$$\frac{\lambda_p}{\lambda_e} = \frac{\frac{h}{m_p v_p}}{\frac{h}{m_e v_e}} \Rightarrow \frac{2}{1} = \frac{m_e v_e}{m_p v_p} = \frac{m_e}{m_p} \times \frac{1}{4}$$

$$\frac{m_e}{m_p} = 8$$

- 22.** In the millikan oil drop experiment radius of drop is  $r = 2 \text{ mm}$  and density  $\rho = 3 \text{ gm/cm}^3$ . If the applied electric field is  $E = 3.55 \times 10^5 \text{ N/C}$ . Find excess electrons.

- (1)  $1.769 \times 10^{10}$       (2)  $1.567 \times 10^{10}$       (3)  $1.769 \times 10^{12}$       (4)  $1.567 \times 10^{12}$


**Ans.** (1)

**Sol.**  $mg = qE$

$$q = \frac{mg}{E}$$

$$N = \frac{mg}{eE} = \frac{3 \times 10^{-3} \times 10 \times \frac{4}{3} \pi \times 8 \times 10^{-9}}{10^{-6} \times 3.55 \times 10^5 \times 1.6 \times 10^{-19}} = 1.769 \times 10^{10}$$

23. A partially filled capacitor has half of its space filled with dielectric of relative permittivity 10. Equivalent capacitance if area of plates is  $100 \text{ m}^2$  and distance between plates is  $10 \text{ m}$  is given as  $x \text{ pF}$ . Find  $x$ ? ( $\epsilon_0 = 8.85 \times 10^{-12}$ )



**Ans.** 161.00

**Sol.**  $C_2 = \frac{\epsilon_0 \times 100}{5} = 20\epsilon_0$


$$C_1 = 10 \times \frac{\epsilon_0 \times 100}{5} = 200\epsilon_0$$

$$C_{eq} = \frac{C_1 C_2}{C_1 + C_2}$$

$$C_{eq} = \frac{4000 \epsilon_0}{220}$$

$$= 160.90 \times 10^{-12} = 161 \text{ pF}$$

24. A ball is released from point A. Evaluate its velocity (m/s) when it reaches to point B (assume frictionless surface):



**Ans.** 10.00

**Sol.**  $mg(5) = \frac{1}{2}mv^2$

$$V = 10 \text{ m/s.}$$

25. Initially a body of mass  $10 \text{ kg}$  is moving along  $x$ -axis with velocity  $10\sqrt{3} \text{ m/s}$ . It collides with another body of mass  $20 \text{ kg}$  and comes to rest. The  $20 \text{ kg}$  mass object disintegrates in 2 parts each of mass  $10 \text{ kg}$ . One part moves along  $y$ -axis with velocity  $10 \text{ m/s}$  and another at  $30^\circ$  with  $x$ -axis. Evaluate the velocity of the object which moves at angle  $30^\circ$  with  $x$ -axis.

**Ans.** 20.00



**Sol.**  $|\vec{v}| = 20 \text{ m/s}$

$$10 \times 10\sqrt{3}\hat{i} = 10 \times 10\hat{j} + 10\vec{v}$$

$$\frac{100\sqrt{3}\hat{i} - 100\hat{j}}{10} = \vec{v}$$

$$\vec{v} = 10\sqrt{3}\hat{i} - 10\hat{j}$$

- 26.** A bullet of mass 0.1 kg initially moving with a velocity 10 m/sec and then passes through a wooden block and comes to rest with uniform deceleration by travelling 50cm. If the force exerted by wooden block on bullet is x newton, then find x.

**Ans.** 10.00

**Sol.**  $v^2 = u^2 + 2as$

$$0 = 100 + 2(-a)\left(\frac{1}{2}\right)$$

$$a = 100 \text{ m/s}^2$$


$$F = ma = (0.1)(100)$$

$$F = 10 \text{ N}$$

- 27.** A capacitor of capacitance  $3 \mu\text{F}$  has charge 30 nC is connected to a resistance of  $5\Omega$ . If current in circuit just after closing the switch is x A. Then x is :

**Ans.** 2.00

**Sol.**



$$q = Qe^{\frac{-t}{RC}}$$

$$I = \frac{Q}{RC} e^{\frac{-t}{RC}}$$

$$I(t=0) = \frac{Q}{RC} = \frac{30}{5 \times 3} = 2\text{A}$$

**28.** Coming soon.

**29.** Coming soon.

**30.** Coming soon.