

JEE
MAIN
2020

NARAYANA'S SENSATIONAL SUCCESS ACROSS INDIA

NARAYANA IIT-JEE (MAIN) 2020 ALL INDIA TOP RANKS IN OPEN CATEGORY



SWAYAM CHUBE
HT No. MR16303069

BELOW
10 **21** RANKS

*All Cat



**HARSHVARDAHAN
AGARWAL**
HT No. HR09300789

BELOW
100 **126** RANKS

*All Cat



SHIVA KRISHNA
HT No. TL01330751

TOTAL
QUALIFIED
FOR JEE-ADV. **16292**

ADMISSIONS OPEN (2020-21)

OUR REGULAR CLASSROOM PROGRAMME

One Year Classroom Program
JEE/NEET-2021
(for students moving from XI to XII)

Two Year Classroom Program
JEE/NEET-2022
(for students moving from X to XI)

Three Year Integrated Classroom Program
JEE/NEET-2023
(for students moving from IX to X)

Four Year Integrated Classroom Program
JEE/NEET-2024
(for students moving from VIII to IX)

FOUNDATION PROGRAMMES
For NTSE, NSEJS, JSTSE,
Olympiads & School/Board Exams
(for students moving to
Class VI, VII, VIII, IX & X)

APEX BATCH
Two years school Integrated
Classroom Program - 2022
For JEE Main & Advance / NEET (for XI Studying Students)
Course Feature - Complete Coverage of CBSE - Regular Classes - Weekly Test & Regular Analysis - Lab Facility
- Motivation & Counseling - Competitive Exam Prep - Ample time for self study

Online Classes for IIT/NEET/Foundation/Olympiads

- Access Recording of Past Classes on n-Learn App
- Online Parent Teacher Meeting
- Personalized Extra Classes & Live Doubt Solving
- Hybrid/Customized Classroom model
- Video Solution of Weekly/Fortnightly Test
- Printed Study Material will be sent by us
- n-Learn App
- Counselling Motivational sessions
- Affordable Fee
- Doubt Classes / Practice Classes
- Provision to Convert from online to regular classroom programme
- Once Classes resume by just paying nominal fee

Online Test

- Micro & Macro Analysis
- Relative performance (All India Ranking)
- Question wise Analysis
- Unlimited Practice Test
- Grand Test

NARAYANA
Digital
Classes
STUDY ONLINE FROM HOME

For Class
7th to 12th+



JEE-MAIN-2021


FEBRUARY ATTEMPT

24.02.21_SHIFT-II

CHEMISTRY

CHEMISTRY

1. Compare the rate of aromatic nucleophilic substitution reaction of the following compounds



- (1) IV > II > III > I (2) III > II > I > IV (3) I > II > III > IV (4) IV > III > II > I

Ans. (4)

Sol. Rate of aromatic nucleophilic substitution reaction depends upon type of halogens and electronic effect of the group present on the ring. Electron withdrawing groups (–I, –M) increases rate of reaction increases.

2. What is S in Buna-S ?

- (1) Sulphure (2) Styrene (3) Rubber (4) Strength


Ans. (2)

Sol. Buna-S is the co-polymer of buta-1,3-diene and styrene

3. Which of the following set of the reagent is used to convert nitrobenzene to 1,3-Dibromobenzene?


- (1) $\text{Br}_2/\text{Fe} \longrightarrow \text{Sn}/\text{HCl} \longrightarrow \text{NaNO}_2/\text{HCl} \longrightarrow \text{CuBr}/\text{HBr}$
 (2)
 (3)
 (4)

Ans. (1)



1,3-Dibromobenzene


4. The reagent used to convert the following is ?



- (1) $\text{NH}_2\text{--NH}_2/\text{C}_2\text{H}_5\text{O}^-\text{Na}^+$ (2) Red P/ Cl_2
 (3) Ni/H_2 (4) NaBH_4

Ans. (1)

Sol.



it is wolff-Kishner reduction of carbonyl compounds.

5. Match the following

Column – I

- (a) Valium
(b) Morphine
(c) Norethindrone
(d) Vitamin B-12

Column – II




- (p) Pernicious anaemia
(q) Analgesic
(r) Tranquilizer
(s) Antifertility

Ans. $a \rightarrow r$; $b \rightarrow q$; $c \rightarrow s$; $d \rightarrow p$ 6. **Statement – I** : BOD is the parameter that can be helpful for survival of aquatic life.**Statement – II** : Optimum value of BOD is 6.5 ppm.

- (1) Statement I is true ,Statement II is false
(2) Statement I is false ,Statement II is true
(3) Statement I , II both are true
(4) Statement I , II both are false



Ans. (1)

7. How many of the following amines can be prepared by Gabriel phthalimide synthesis ?







Ans. (3)

Sol. Only aliphatic amines can be prepared by Gabriel phthalimide synthesis.

8. Which of the following compound cannot be prepared by the reaction of alkyne with $\text{HgSO}_4/\text{dil. H}_2\text{SO}_4$?

Ans. (2)

9. Diazonium salt of which of the following will give coloured dye on reaction with β -Naphthol in NaOH 


Ans. (4)

Sol. Only aromatic Primary amines will gives Dye test.

10. The correct bond angle & shape of I_3^- is

- (1) Linear & 180° (2) Trigonal pyramidal & 120°
 (3) V-shape & 120° (4) T-shape & $109^\circ 28'$

Ans. (1)



Linear shape $\angle \text{I}-\text{I}-\text{I} = 180^\circ$

11. Correct statements

(a) $\text{K.E.} \propto \frac{Z^2}{n^2}$

(b) $(nv) \propto Z^2$

(c) $\text{Frequency} \propto \frac{Z^3}{n^3}$

(d) $\text{Electrostatic force} \propto \frac{Z^3}{n^4}$

- (1) a & d are correct (2) a & b are correct
 (3) b & c are correct (4) b & d are correct

Ans. (1)

12. Which of the following is incorrect?

- (1) Cr_2O_3 is Amphoteric (2) RuO_4 is oxidising agent
 (3) VOSO_4 is reducing agent (4) Ruby appears due to presence of Co^{3+}

Ans. (4)

13. Which of the following order of melting point is correct

- (1) $\text{LiF} > \text{LiCl}$, $\text{NaCl} > \text{MgO}$ (2) $\text{LiF} < \text{LiCl}$, $\text{NaCl} > \text{MgO}$
 (3) $\text{LiF} > \text{LiCl}$, $\text{NaCl} < \text{MgO}$ (4) $\text{LiF} < \text{LiCl}$, $\text{NaCl} > \text{MgO}$

Ans. (3)

Sol. Lattice energy $\propto \frac{1}{r^+ + r^-}$

$$\propto \frac{1}{r^+ + r^-}$$

LiF LiCl

Size $\text{F}^- < \text{Cl}^-$ [charge are same]


Lattice energy $\boxed{\text{LiF} > \text{LiCl}}$



Lattice energy $\text{MgO} > \text{NaCl}$

Charge dominate over size

14. Spin only magnetic moment of the following complexes



(1) 4.9, 0, 1.76 BM

(2) 5.9, 1.73 BM

(3) 1.73, 2.83, 0 BM

(4) 2.83, 6.9, 0 BM

Ans. (1)

Sol. $[\text{FeCl}_4]^{2-}$ Contain Fe^{+2} in tetrahedral complex. Its configuration is $e_g^{2,1} t_{2g}^{1,1,1}$ it have 4 unpaired electron in $[\text{Co}(\text{ox})_3]^{3-}$ Co^{+3} have configuration $t_{2y}^{2,2,2} e_g^{0,0}$ MnO_4^{2-} have Mn in +6 oxidation state and configuration of Mn is $e_g^{1,0} t_{2g}^{0,0,0}$

15. α -sulphur, β -Sulphur, $\text{S}_2 \rightarrow$ find how many are paramagnetic

Ans. (1)

Sol. In S_2 , like O_2 two unpaired electron are present, α & β - sulphur have S_8 ring which are diamagnetic

16. Which of the following can be used for clotting of blood efficiently?

(1) NaHCO_3


(2) FeCl_3

(3) FeSO_4

(4) $\text{Mg}(\text{HCO}_3)_2$

Ans. (2)

Sol. Blood is a negative charged Sol. Therefore according hardy-Schulz rule Fe^{+3} cation have highest coagulation power. Therefore FeCl_3 can be used for clotting of blood efficiently.



given that

$$G_m^\circ (\text{C}_2\text{H}_2) = 2.4 \times 10^5 \text{ J}$$

$$G_m^\circ (\text{C}_6\text{H}_6) = -1.4 \times 10^5 \text{ J}$$

Find $\log_{10} k$ at 25°C

Ans. (150.72)

Sol. $\Delta G^\circ = (G_M^\circ)_{C_6H_6} - 3(G_M^\circ)_{C_2H_2}$

$$= -1.4 \times 10^5 - 3 \times 2.4 \times 10^5$$

$$= -8.6 \times 10^5 \text{ Joule}$$


$$-2.303RT \log_{10} k = -8.6 \times 10^5$$

$$-2.303 \times 8.314 \times 298 \log_{10} k = -8.6 \times 10^5$$

$$\log_{10} k = 150.72$$

- 18.** 1.86 gm of aniline is converted into acetanilide with 90% efficiency. Mass of acetanilide formed is $[X] \times 10^{-2}$ gm

Ans. 243×10^{-2}



1.86 g

Molar mass = 93

Molar mass = 135

* 93 g aniline produces 135 g acetanilide

$$1.86 \text{ g aniline produces } \frac{135 \times 1.86}{93} = 2.70 \text{ g}$$

* At 90% efficiency of reaction it produces $= \frac{2.70 \times 90}{100} = 2.43 \text{ g}$

Ans. 243×10^{-2}

- 19.** Freezing point of C_6H_6 (ℓ) is 5.5°C . If 10g of C_4H_{10} is mixed with 200g of C_6H_6 (ℓ). Calculate freezing point of solution $k_f = 5.12^\circ\text{C/m}$.

Ans. (1.09°C)

Sol. $\Delta T_f = k_f \times m$

$$= 5.12 \times \frac{10}{58} \times \frac{1000}{200} = 4.41^\circ\text{C}$$

$$\Delta T_f = (T_f)_{\text{Solvent}} - (T_f)_{\text{Solution}}$$

$$4.41^\circ\text{C} = 5.5 - (T_f)_{\text{Solution}}$$

$$(T_f)_{\text{Solution}} = 5.5 - 4.41 = 1.09^\circ\text{C}$$

20. De-broglie's wavelength of a proton and an α -particle is same. Calculate ratio of their velocities

Ans. (4)

Sol. $\lambda_p = \lambda_\alpha$

$$\frac{h}{m_p v_p} = \frac{h}{m_\alpha v_\alpha}$$

$$\frac{v_p}{v_\alpha} = \frac{m_\alpha}{m_p} \quad \because m_\alpha = 4 m_p$$

$$\frac{v_p}{v_\alpha} = \frac{4m_p}{m_p} = 4$$

Ans. 4

21. If $[H^+]$ changed from 1M to 10^{-4} M

Find change in electrode potential $E^\circ_{MnO_4^-/Mn^{+2}}, \left(\frac{RT}{F} = 0.059\right)$

[Assume $[MnO_4^-] = [Mn^{+2}] = 1M$]

Ans. 0.3776 V

Sol. $5e^- + 8H^+ + \underset{1M}{MnO_4^-} \longrightarrow \underset{1M}{Mn^{+2}} + 4H_2O$

$$E_1 = E^\circ - \frac{0.59}{5} \log_{10} \left[\frac{1}{[H^+]^8} \times \frac{[Mn^{+2}]}{[MnO_4^-]} \right]$$

$$= E^\circ - \frac{0.059}{5} \log_{10} \left[\frac{1}{(1)^8} \right] = E^\circ$$

$$E_2 = E^\circ - \frac{0.059}{5} \log_{10} \left[\frac{1}{(10^{-4})^8} \times \frac{[Mn^{+2}]}{[MnO_4^-]} \right]$$

$$= E^\circ - \frac{0.059}{5} \log_{10} [10^{32}]$$


$$= E^\circ - \frac{0.059}{5} \times 32$$

$$E_1 - E_2 = E^\circ - E^\circ + \frac{0.059}{5} \times 32$$

$$= 0.3776 \text{ V}$$

22. V ml of a hydrocarbon C_xH_y requires 6V ml of oxygen for complete combustion & forms 4V ml of CO_2 . Determine y

Ans. 8



Volume-Volume V 6V 4V

Analysis

$$\frac{V_{C_xH_y}}{1} = \frac{V_{CO_2}}{x}$$

$$\frac{v}{1} = \frac{4v}{x} \quad x = 4$$

$$\frac{v_{C_xH_y}}{1} = \frac{V_{O_2}}{x + \frac{y}{4}}$$

$$\frac{V}{1} = \frac{6V}{x + \frac{y}{4}}$$

$$x + \frac{y}{4} = 6$$

$$4 + \frac{y}{4} = 6$$

$$\frac{y}{4} = 2$$

$$y = 8$$

Formula C_4H_8

23. Sucrose $\xrightarrow{\text{I order}}$ Glucose + Fructose

$$t_{1/2} = 3.33 \text{ hour}$$

f = fraction remaining of sucrose at 9 hour.

Find out value of $100 \times \log \left(\frac{1}{f} \right)$

$$[\log_{10} 2 = 0.3]$$

Ans. (81)

Sol. $f = \frac{1}{2^n}$ $n = \frac{\text{time given}}{t_{1/2}} = \frac{9 \text{ hr}}{3.33 \text{ hr}} = 2.7$

$$= \frac{1}{2^{2.7}}$$

$$\log \frac{1}{f} = \log 2^{2.7} = 2.7 \times 0.3 = 0.81$$

$$100 \times \log_{10} \left(\frac{1}{f} \right) = 100 \times 0.81 = 81$$

Ans. 81

- 24.** Determine volume occupied by 4.75g acetylene gas at 740 mmHg pressure & 50°C temperature
R = 0.0826 Latm/mol k (in L)

Ans. (5)

Sol.
$$V = \frac{nRT}{P} = \frac{\left(\frac{4.75}{26} \right) \times 0.0826 \times 323}{\left(\frac{740}{760} \right)} \approx 5\text{L}$$